

# The Interest in Kaonic-Deuterium Atoms

S. Wycech\*

*Soltan Institute for Nuclear Studies, Warsaw, Poland*

## Abstract

K meson-deuteron scattering lengths, atomic level shifts and widths are calculated. These quantities depend on controversial kaon-nucleon scattering amplitudes extrapolated by few MeV below the  $KN$  threshold. Experimental consequences of several  $KN$  interaction models are studied.

---

\*e-mail "wycech@fuw.edu.pl"

## I. INTRODUCTION

Few-body interactions of the  $K^-$  meson may complement our knowledge on the kaon-nucleon interaction. One point of interest in this field is the old question of the nature of  $\Lambda(1405)$  state interpreted as an elementary particle [1] or a  $KN$  quasi-bound state [2]. This uncertainty is aggravated by a serious discrepancy between the value of  $Kp$  scattering length obtained in scattering experiments [3] and the value extracted from the  $1S$  level shift in kaonic-hydrogen atoms [4], [5], [6], [7]. Of related interest is the possibility of  $K$ -nucleus quasi-bound states [11], the question of kaon condensation and its interesting astrophysical consequences [12]. Another controversial, misunderstood but apparently related result is the anomalously large  $2P$ -state width and shift detected in the  $K^-$ -He atoms [8], [10], [9]. The helium measurements indicate possible existence of a nuclear bound state in the  $K^- \alpha$  system.

These unsolved problems have been dormant for a decade, due to lack of intense low energy kaonic beams. Such a beam is now coming into operation at the DAΦNE facility and the kaonic atom program DEAR is being installed there [13]. Another kaonic atom program is running now at KEK [7]. The purpose of this letter is to study the physical interest and chances of the kaonic deuterium X-ray measurements.

The  $K^-$  deuteron system is in general difficult to describe because of numerous inelastic decay channels. However, to study the kaonic-deuterium atoms one needs to know only the  $KD$  elastic scattering amplitude at threshold. This could be obtained from the elastic  $KN$  amplitudes by summation of the multiple scattering series. Such a description involves elementary  $KN$  scattering amplitudes extrapolated into the subthreshold region. Such extrapolation is necessary since the nucleons are bound. In addition, the recoil energy of the  $KN$  pair reduces the energy available in its C.M. system by several MeV. Thus in the  $KD$  atoms the energies allowed in the  $KN$  subsystems are located half-way between the  $KN$  threshold and the  $\Lambda(1405)$ . On the other side, the  $K^-$ -He atoms involve  $KN$  energies down in the interesting  $\Lambda(1405)$  region as the nucleons are bound more strongly in the  $\alpha$

particle. The price to pay is the difficulty in a reliable description of the  $K\alpha$  system. It is the deuteron that offers a good compromise between the requirements of theories and the atomic experiments.

In this letter a simple formula for the  $KD$  scattering length, volume etc. is given and related shifts and widths of atomic levels are calculated. This formula is expressed by certain averages of the  $Kn$  and  $Kp$   $S$ -wave scattering amplitudes below the thresholds. These averages are our free parameters and are taken from theoretical models or phenomenological scattering analyses. It turns out that it is the widths of atomic states that are most sensitive to these input parameters. The  $1S$  level shifts display rather limited sensitivity to the real amplitudes and are dominated by the absorptive parts of  $KN$  amplitudes. The results indicate a good chance for a successful measurement within the DEAR project.

## II. RELATION OF LEVEL SHIFTS TO SCATTERING AMPLITUDES

The atomic level shifts  $\Delta E$  induced by nuclear interactions may be related to the scattering lengths of the orbital meson on atomic nuclei. These relations are very simple when the atomic Bohr radius  $B$  is larger than the lengths characteristic for nuclear interactions : the nuclear radius  $R_D$  and the meson-nucleus scattering length  $A_0$ . For the  $S$  waves such a relation

$$\Delta E_{nS} = E_{nS} - \epsilon_{nS} = \frac{2\pi}{\mu} |\psi_n(0)|^2 A_0 (1 - \lambda A_0/B) \quad (1)$$

is known as the Trueman formula [14]. It is accurate to a second order in  $A_0/B$ , good enough for the  $KD$  system. The electromagnetic energy  $\epsilon_{nS}$  in Eq.(1) is composed of the Bohr's atom energy  $\epsilon_{nS}^o = -\mu(\alpha)^2/2n^2$  corrected for relativity and effects of the deuteron electric polarisability. The Bohr radius is given by  $= 1/\alpha\mu$  where  $\mu$  is the reduced mass. In the  $1S$  state one has  $\lambda = 3.154, B = 69.2fm, A \approx 1fm$  and the second order term in Eq.(1) constitutes a few percent correction. Such corrections are negligible in higher angular momentum states. For these states a simpler linear relation

$$\Delta E_{nL} = \epsilon_{nL}^o \frac{4}{n} \Pi_{i=1}^L \left( \frac{1}{i^2} - \frac{1}{n^2} \right) A_L / B^{2L+1} = -\Theta_{nL} \frac{A_L}{\mu B^{2L+1}} \quad (2)$$

due to Lambert [15] is sufficient. We shall be interested in the lowest circular states  $1S, 2P, 3D$ . For these the overlap factors  $\Theta$  equal correspondingly  $2, 3/16, 60/(81)^2$ .

Our problem is to calculate the  $KD$  scattering length  $A_0$ , scattering volume  $A_1$  etc. These quantities are obtained in the next section by a trick which sums the  $KD$  multiple scattering series. This method was tested with success in the  $\eta D$  calculations where the details of the method are explained [17]. In this way the kaon-deuteron scattering lengths are expressed in terms of deuteron radius, properties of  $NN$  interactions and the  $KN$  scattering amplitudes. The latter are complex due to  $KN$  coupling with the open  $\Sigma\pi, \Lambda\pi$  channels. The absorptive parts of these amplitudes contribute to absorptive parts of the  $KD$  scattering lengths. The latter determine the widths of atomic  $KD$  states.

### III. A FORMULA FOR THE KAON-DEUTERON SCATTERING MATRIX

The purpose of this section is to exploit a simple formula that relates the meson-deuteron scattering lengths to the meson-nucleon subthreshold scattering amplitudes. The  $KD$  lengths are found by a summation of the multiple scattering series and are expressed in terms of few basic multiple scattering integrals. Although we are interested in all partial waves in the  $KD$  system only the  $S$  wave interactions in the  $KN$  system are considered. To keep the presentation simple, the impulse approximation is discussed in some detail but higher order terms and the summation of the series is presented rather schematically. For details we refer to a similar calculation of  $\eta$ -deuteron scattering length [17].

Let us begin with a multiple scattering expansion that follows from the three-body Faddeev equations for a meson interacting with a pair of nucleons labeled 1,2. For the situation of meson-deuteron scattering below the deuteron breakup, the series for the kaon-deuteron  $T_{KD}$ -matrix is:

$$T_{KD} = t_1 + t_2 + t_1 G_0 t_2 + t_2 G_0 t_1 + t_2 G_0 t_1 G_0 t_2 + t_1 G_0 t_2 G_0 t_1$$

$$+ (t_1 + t_2)G_{NN}(t_1 + t_2) + (t_1 + t_2)G_{NN}(t_1 + t_2)G_{NN}(t_1 + t_2) + \dots \quad (3)$$

where  $t_i$  are meson-nucleon scattering matrices,  $G_0$  is the free three-body propagator and  $G_{NN} = G_0 T_{NN} G_0$  is that part of the three-body propagator which contains the nucleon-nucleon scattering matrix  $T_{NN}$ . This expansion is performed in the momentum space and appropriate integrations over the intermediate momenta are understood.

The scattering amplitude is determined by an average

$$< \psi_D \psi_K^L | T_{KD}(E) | \psi_D \psi_K^L > \equiv < L | T_{KD}(E) | L > \quad (4)$$

where  $\psi_D$  is the deuteron and  $\psi_K^L$  is the free-meson wave function for an  $L$ -th partial wave in the meson-deuteron system. We define the  $KD$  scattering lengths as the limit for  $p \rightarrow 0$  of the expression

$$A_{KD}(L) = -(2\pi)^2 m_{KD} < L | T_{KD}/p^{2L} | L > \quad (5)$$

where  $p$  is a relative meson-deuteron momentum and  $m_{KD}$  is the corresponding reduced mass. The normalization in Eq.5 is related to the normalisation of  $t_{KN}$

$$t_{KN}(E) = -\frac{a_{KN}(E)}{(2\pi)^2 \mu_{KN}} \quad (6)$$

where  $a_{KN}(E)$  is the  $S$  wave scattering amplitude in the  $KN$  system normalised to the scattering length at the  $KN$  threshold. Here,  $\mu_{KN}$  is another reduced mass.

Calculations of the multiple scattering integrals are straightforward although tedious and the formulae are lengthy. For an illustration we reproduce the leading impulse approximation lengths for zero range  $KN$  forces. Actual calculations are performed with a finite force range generated by a separable Yamaguchi potential of 600 MeV inverse range .

The leading term  $< L | T_{KD}^0 | L > = < L | t_1 + t_2 | L >$  is the impulse approximation. This term yields the impulse  $KD$  scattering lengths

$$A_{KD}^0(L) = \frac{\mu_{KD}}{\mu_{KN}} (\bar{a}_{Kp} + \bar{a}_{Kn}) \frac{< (r/2)^{2L} >_D}{(2L+1)!!^2} \quad (7)$$

where  $< (r/2)^{2L} >_D$  is the  $2L$ -th radial moment of the deuteron related to its center and

$$\bar{a} = \int a(-E_D - \frac{p^2}{2m_{N,KN}}) |\tilde{\phi}_D^L(p)|^2 d\vec{p} \int |\tilde{\phi}_D^L(p)|^2 d\vec{p} \quad (8)$$

are the scattering matrices averaged over some energy region, generated by the recoil of the spectator nucleon. The range of the latter is given by a Bessel transform of the deuteron wave function

$$\tilde{\phi}(p) = \int \psi_D(r) j_L(pr/2) p^2 dp \quad (9)$$

The energies involved in Eq.(8) extend down to the unphysical subthreshold region. The recoil energy region given by the Bessel transform depends on the partial wave  $L$ . The relevant distributions peak around energies of -12,-7 and -5 MeV for  $L = 0, 1, 2$  correspondingly. Therefore, a model is required for this extrapolation and several possibilities are discussed later.

Now a partial summation of the series (3) for  $\langle T \rangle$  is obtained by

$$\langle T_{KD}^1 \rangle = \frac{\langle T_{KD}^0 \rangle}{1 - \Omega_1 - \Sigma_1} \quad (10)$$

with

$$\Omega_1 = \frac{\langle t_1 G_0 t_2 + t_2 G_0 t_1 \rangle}{\langle T_{KD}^0 \rangle} \quad (11)$$

$$\Sigma_1 = \frac{\langle T_{KD}^0 G_{NN} T_{KD}^0 \rangle}{\langle T_{KD}^0 \rangle}. \quad (12)$$

This partial sum is very close to the formula for meson scattering on two fixed nucleons [16]. It contains terms of the order  $\xi = \langle t \rangle / R_D$  in the denominator. The parameter  $\xi$  does not need to be small for the success of Eq.(10), which works also for  $|\xi| > 1$ , when the multiple scattering is divergent. In the  $K - D$  case  $|\xi|$  falls into 0.3-.5 range. Corrections of higher orders of  $\xi$  in the denominator of Eq.(10) may be obtained by comparing higher orders in Eq.(3) with a series expansion of Eq.(10) with respect to  $\Omega_1$  and  $\Sigma_1$ . In this way the second approximation is obtained

$$\langle T_{KD}^2 \rangle = \frac{\langle T_{KD}^0 \rangle}{1 - \Omega_1 - \Sigma_1 [\Omega_2 - (\Omega_1)^2] - [\Sigma_2 - (\Sigma_1)^2] - [\Delta_2 - \Omega_1 \Sigma_1]} \quad (13)$$

where

$$\Omega_2 = \frac{\langle t_1 G_0 t_2 G_0 t_1 + t_2 G_0 t_1 G_0 t_2 \rangle}{\langle T_{KD}^0 \rangle} \quad (14)$$

and

$$\Sigma_2 = \frac{\langle T_{KD}^0 G_{NN} T_{KD}^0 G_{NN} T_{KD}^0 \rangle}{\langle T_{KD}^0 \rangle}. \quad (15)$$

$$\Delta_2 = 2 \frac{\langle T_{KD}^0 G_{NN} (t_1 G_0 t_2 + t_2 G_0 t_1) \rangle}{\langle T_{KD}^0 \rangle} \quad (16)$$

This procedure may be continued but the main advantage of this order is a strong cancellation in the  $\Sigma_2 - \Sigma_1^2$  term (and indeed also in the higher order  $\Sigma_n$  terms).

The convergence of our expansion is shown in Table 1 in the case of  $S$  wave scattering length. It seems quite satisfactory and a percent precision is reached. The details of the calculations may be found in Ref. [17] where this formalism is used to describe the  $\eta$  deuteron scattering.

For completeness, the scattering length should contain also an *inner* Coulomb correction. A quick estimate follows from a boundary condition model that generates a relation  $1/A_c = 1/A - 2/B(\ln(2R_D/B) + 1.154)$  where  $A_c$  is the Coulomb corrected length. The change is on a percent level and will be neglected in the following estimates.

## IV. RESULTS

The meson-deuteron scattering length  $A_{KD}$  is expressed in terms of "effective  $Kp$  and  $Kn$  scattering lengths"  $\bar{a}_{KN}$ , which are some averages of the  $KN$  scattering matrices extrapolated below the threshold. These are input parameters taken from several analyses of the combined  $\pi\sigma, \pi\Lambda, KN$  coupled channels data. A collection of characteristic numbers is given in Table 2.

In Table 3 one finds values of the atomic level shifts and half-widths for the  $KN$  effective scattering lengths given in Table 2. The resulting  $S$  wave length  $A_{KD}$  and related level

width may vary by an order of magnitude, reflecting the nearby  $KD$  quasibound state that may arise at threshold for  $\Re a_{KN}$  close to 1 fm. It is clear that such a possibility exists for one of the models discussed [23], [20]. Other models are rather distant from this situation.

The effective lengths were grouped into two sets. Those of Refs. [18], [2], [19], [22] fit the scattering results and disagree with the  $1S$  kaonic hydrogen level data. The differences of the  $KD$  level shifts and widths are rather moderate, still some of those, in particular the  $2P$  level widths could be discernible by the atomic cascade measurements. The differences involved in these models are partly related to the subthreshold extrapolations which are known to depend on details of the interaction. Other differences are due to different nonresonant  $I = 1$  amplitudes that weight heavily in the deuteron case.

Other, less conventional, models tend to describe the hydrogen level shift and width. Those [23], [21], [20] generate significantly smaller  $KN$  absorptive amplitudes. As a consequence the widths of  $2P, 3D$  states in the deuterium are also much smaller. Another interesting physical consequence of some of these models is a near-possibility of a nuclear  $S$  wave  $KD$  state. Such a state is almost-generated by the attractive  $Kn$  interactions involved in models of Refs. [23], [20]. Proximity of such a state is reflected in the enhanced widths of the  $1S$  kaonic-deuterium atomic level indicated in Table 3.

Two conclusions follow from this paper:

- a) Different parametrisations and models of kaon-nucleon interactions generate sizable differences in the kaonic-deuteron level shifts and lifetimes.
- b) Two groups of models, that reproduce either the scattering or the hydrogen data, generate very different  $2P, 3D$  deuteron widths. These two situations should be discernible by the atomic cascade in the kaonic deuterium even if the shape of  $2P \rightarrow 1S$  line turns out difficult to resolve.



## TABLES

TABLE I. Convergence of the  $A_{KD}(L=0)$  expansion. Last line is a third order formula, not included in the text.  $\bar{a}_{Kn} = -0.1 + i0.5, \bar{a}_{Kp} = -1. + i1.5$

|            |                 |                |
|------------|-----------------|----------------|
| $A_{KD}^0$ | $-1.39 + i2.42$ | <i>Eq.(7)</i>  |
| $A_{KD}^1$ | $-1.38 + i0.74$ | <i>Eq.(10)</i> |
| $A_{KD}^2$ | $-1.23 + i0.95$ | <i>Eq.(13)</i> |
| $A_{KD}^3$ | $-1.24 + i0.95$ | <i>Order3</i>  |

TABLE II. Kaon-nucleon "effective lengths" in  $fm$  units. These numbers have been calculated from figures in the referred papers, hence contain some numerical uncertainties and have been rounded correspondingly.

|                    | $a_{\bar{K}p}$ | $a_{\bar{K}n}$ |
|--------------------|----------------|----------------|
| Martin [2]         | $-1.0 + i1.5$  | $-0.1 + i0.5$  |
| Chao [18]          | $-0.7 + i1.8$  | $0.4 + i0.7$   |
| Ha [3] [22]        | $-1.0 + i2.0$  | $-0.1 + i0.7$  |
| Waas [19]          | $-0.5 + i2.2$  | $0.8 + i0.5$   |
| Tanaka [23]        | $0.0 + i0.7$   | $1.1 + i0.5$   |
| Kumar [21]         | $-1.0 + i0.1$  | $0.4 + i0.7$   |
| Schnick [20], [23] | $0.6 + i0.5$   | $1.1 + i0.2$   |

TABLE III. Level shifts and half-widths of the  $1S, 2P, 3D$  states in kaonic-deuterium. The units are  $KeV, meV, 10^{-2}\mu eV$  correspondingly.

|                    | $\Delta E - i\Gamma/2_{1S}$ | $\Delta E - i\Gamma/2_{2P}$ | $-i\Gamma/2_{3D}$ |
|--------------------|-----------------------------|-----------------------------|-------------------|
| Martin [2]         | 0.76 - i0.64                | 6.5 - i12                   | - i6.9            |
| Chao [18]          | 0.74 - i0.78                | 1.4 - i15                   | - i8.5            |
| Ha [3] [22]        | -0.81 - i0.75               | 5.7 - i16                   | - i9.3            |
| Waas [19]          | 0.83 - i0.87                | -2.5 - i16                  | - i9.4            |
| Tanaka [23]        | 0.29 - i1.22                | -6.7 - i6.9                 | - i4.2            |
| Kumar [21]         | 0.43 - i0.19                | 3.8 - i4.0                  | - i2.3            |
| Schnick [20], [23] | 0.58 - i2.87                | -9.8 - i3.7                 | - i2.4            |

## REFERENCES

- [1] R.H. Dalitz and J.G. McGinley, in Low and Interm. Energy KN Physics ed. E. Ferrari and G.Violini (Dodrecht:Riedel)(1980)381
- [2] A.D. Martin Phys. Lett. B63(1976) 266,Nucl.Phys.B179(1980)33
- [3] J.Ciborowski et al, J.Phys. G9(1982)13
- [4] J.Davies et al Phys.Let B83(1979)55
- [5] M.Izycki et al Zeit. Phys. A297(1980)11
- [6] P.M Bird et al Nucl.Phys. A404(1983)482
- [7] M. Iwasaki et al , Nucl. Phys. A585(1995)239c
- [8] C.F.Wiegand and R.H.Pehl, Phys.Rev.Let 27(1971)1410
- [9] C. Batty et al, Nucl.Phys.A326(1983)455
- [10] S. Baird et al Nucl.Phys. A392(1983)297
- [11] S. Wycech, Nucl.Phys. A450(1986)399c
- [12] G.E. Brown, Nucl.Phys. A574(1994)217c
- [13] The DEAR proposal at DA  $\Phi$  NE, INFN Frascati, Rome
- [14] T.L. Trueman, Nucl. Phys. 26(1961)57
- [15] E.Lambert,Helv. Phys. Acta 43(1970)713
- [16] K.A. Brueckner, Phys.Rev.90(1953)715
- [17] A.M.Green,J.A.Niskanen,and S.Wycech, Phys. Rev C 54,(1996)1970.
- [18] Y.A. Chao et al, Nucl.Phys. B56(1973)46.
- [19] T.Waas, N.Kaiser and W.Weise, Phys.Lett. B365(1996)12.

- [20] J. Schnick and R.H.Landau, Phys.Rev.Lett.58(1987)1719.
- [21] K.S.Kumar and Y.Nogami, Phys. Rev D21,(1980)1834.
- [22] G. He and R.H.Landau, Phys.Rev C48,(1993)3047.
- [23] K.Tanaka and A.Suzuki, Phys.Rev C45,(1992)2068.